



Localized excitations in Debye crystals: a survey of theoretical results

Ioannis Kourakis^{1,2}, Vassilios Koukoulouyannis³ and Padma Kant Shukla²

¹ Centre for Plasma Physics (CPP), Queen's University Belfast, BT7 1 NN Northern Ireland, UK; Email: i.kourakis@qub.ac.uk, www.kourakis.eu

² Institut für Theoretische Physik IV, Ruhr-Universität Bochum, Germany; ³ School of Physics, Theoretical Mechanics, Aristotle University of Thessaloniki, Greece



1. Introduction

Nonlinear processes in *strongly coupled* dusty plasmas (DP) have attracted theoretical interest recently, motivated by recent experiments. Dust (quasi-)lattices (DL) (2D or 3D) are typically formed in the sheath region above the negative electrode in discharge experiments, horizontally suspended at a levitated equilibrium position, at $z = z_0$, where gravity and electric (and/or magnetic) forces balance. Appropriate trapping potentials have also enabled the realization of 1D lattices, dominated by electrostatic interactions.




FIG. 10. Experimental configuration for forming a linear dust chain above a long rectangular box on a negatively biased mesh electrode. (from [1a])

The linear regime of low-frequency oscillations in DP crystals, in the longitudinal (acoustic mode) and transverse (in-plane, shear acoustic mode and vertical, off-plane optical mode) direction(s), is now quite well understood. However, the *nonlinear* (NL) behaviour of DP crystals is little explored, and has lately attracted experimental [1-3] and theoretical [4-7] interest. We have recently considered the coupling among the horizontal ($\sim \hat{x}$) and vertical (off-plane, $\sim \hat{z}$) degrees of freedom in dust mono-layers; a set of NL equations for coupled longitudinal and transverse dust lattice (LDL, TDL) motion was thus derived [4].

Here, we review the nonlinear dust grain excitations which may occur in a DP crystal (assumed quasi-one-dimensional and infinite, composed from identical grains, of equilibrium charge q and mass M , located at $x_n = nr_0$, $n \in \mathcal{N}$). Damping is omitted here.

2. Transverse envelope structures (continuum)

Taking into account the intrinsic nonlinearity of the sheath electric (and/or magnetic) potential, the vertical (off-plane) n -th grain displacement $\delta z_n = z_n - z_0$ in a dust crystal (where $n = \dots, -1, 0, 1, 2, \dots$), obeys the equation

$$\frac{d^2 \delta z_n}{dt^2} + \nu \frac{d(\delta z_n)}{dt} + \omega_{T,0}^2 (\delta z_{n+1} + \delta z_{n-1} - 2\delta z_n) + \omega_g^2 \delta z_n + \alpha (\delta z_n)^2 + \beta (\delta z_n)^3 = 0. \quad (1)$$

(where coupling anharmonicity and second+ neighbor interactions are omitted)

The characteristic frequency


$$\omega_{T,0} = [-qU'(r_0)/(Mr_0)]^{1/2}$$

is related to the (electrostatic) interaction potential; for a Debye-Hückel potential: $U_D(r) = (q/r) e^{-r/\lambda_D}$, one has

$$\omega_{0,D}^2 = \omega_{DL}^2 \exp(-\kappa) (1 + \kappa)/\kappa^3$$

$\omega_{DL} = [q^2/(M\lambda_D^3)]^{1/2}$ is the characteristic dust-lattice frequency; λ_D is the Debye length; $\kappa = r_0/\lambda_D$ is the DP lattice parameter. The on-site electric potential near equilibrium ($z = z_0$) reads

$$\Phi(z) \approx \Phi(z_0) + M[\omega_g^2 \delta z_n^2/2 + \alpha (\delta z_n)^3/3 + \beta (\delta z_n)^4/4] + \dots$$




(G. Sorasio et al., 2002)

(in account of the electric and/or magnetic field inhomogeneity and charge variations), which is related to the overall vertical force

$$F(z) = F_{el/m}(z) - Mg \equiv -\partial\Phi(z)/\partial z.$$

Linear excitations, viz. $\delta z_n \sim \cos \phi_n$ (here $\phi_n = nkr_0 - \omega t$; k and ω are the wavenumber and frequency; damping is neglected) obey the *optic-like discrete backward wave* [1] dispersion relation

$$\omega^2 = \omega_g^2 - 4\omega_{T,0}^2 \sin^2(kr_0/2) \equiv \omega_T^2. \quad (2)$$



(from [1c])

A multiple scale technique for a *continuum* wavepacket gives [5]:

$$\delta z_n \approx \epsilon (A e^{i\phi_n} + \text{c.c.}) + \epsilon^2 [w_0^{(2)} + (w_2^{(2)} e^{2i\phi_n} + \text{c.c.})] + \dots$$

where $w_0^{(2)} \sim |A|^2$, $w_2^{(2)} \sim A^2$.

The amplitude A obeys the *nonlinear Schrödinger equation*:

$$i \frac{\partial A}{\partial T} + P \frac{\partial^2 A}{\partial X^2} + Q |A|^2 A = 0, \quad (3)$$

where $\{X, T\}$ are the *slow* variables $\{\epsilon(x - v_g t), \epsilon^2 t\}$.

The *dispersion coefficient* $P_T = \omega_T''(k)/2$ takes negative (positive) values for low (high) k .

The *nonlinearity coefficient* $Q = [10\alpha^2/(3\omega_g^2) - 3\beta]/2\omega_T$ is positive for *all* known experimental values of α, β [3].

For small wavenumbers k (where $PQ < 0$), TDLWs will be modulationally stable, and may propagate in the form of dark/grey envelope excitations (*hole solitons* or *voids*) [5].

For larger k , *modulational instability* may lead to the formation of bright (*pulse*) envelope solitons.

Exact expressions for these excitations can be found in [5].




Fig. Envelope solitons of the (a, b) bright type; (c, d) dark (black/grey) type.

3. Transverse Intrinsic Localized Modes (ILMs) – Discrete Breathers (DBs)

ILMs, i.e. highly localized *Discrete Breather* (DB) and *multi-breather*-type few-site vibrations have recently received increased interest among researchers in solid state physics, due to their omnipresence in periodic lattices and remarkable physical properties [6]. Dusty plasma DB excitations were shown to occur in transverse DL motion [7-10] from first principles (figure from [9]).




FIG. 5. (Color online) Time evolution of a 1:1 two-breather for $\epsilon=0.016$. FIG. 6. (Color online) Time evolution of a 2:3 two-breather for $\epsilon=0.003$.

The existence of such DB structures at a frequency ω_{DB} generally requires the *non-resonance condition*

$$n\omega_{DB} \neq \omega(k) \quad \forall n \in \mathcal{N}$$

which is, remarkably, satisfied in all known TDLW experiments [1]. The existence of DBs in 2D (hexagonal) dusty plasma structures is now under investigation [10].

4. Longitudinal envelope excitations

The NL longitudinal equation of motion ($\delta x_n = x_n - nr_0$) reads:

$$\frac{d^2(\delta x_n)}{dt^2} + \nu \frac{d(\delta x_n)}{dt} = \omega_{0,L}^2 (\delta x_{n+1} + \delta x_{n-1} - 2\delta x_n) - a_{20} [(\delta x_{n+1} - \delta x_n)^2 - (\delta x_n - \delta x_{n-1})^2] + a_{30} [(\delta x_{n+1} - \delta x_n)^3 - (\delta x_n - \delta x_{n-1})^3], \quad (4)$$

where the characteristic frequency is given by

$$\omega_{0,L}^2 = [U'''(r_0)/M] = 2\omega_{DL}^2 \exp(-\kappa) (1 + \kappa + \kappa^2/2)/\kappa^3$$

for Debye interactions.

The resulting *acoustic* linear mode⁴ obeys

$$\omega^2 = 4\omega_{L,0}^2 \sin^2(kr_0/2) \equiv \omega_L^2.$$

One now obtains (to lowest order $\sim \epsilon$)

$$\delta x_n \approx \epsilon [u_0^{(1)} + (u_1^{(1)} e^{i\phi_n} + \text{c.c.})] + \epsilon^2 (u_2^{(2)} e^{2i\phi_n} + \text{c.c.}) + \dots,$$

where $u_1^{(1)}$ obey [11]

$$i \frac{\partial u_1^{(1)}}{\partial T} + P_L \frac{\partial^2 u_1^{(1)}}{\partial X^2} + Q_0 |u_1^{(1)}|^2 u_1^{(1)} + \frac{p_0 k^2}{2\omega_L} u_1^{(1)} \frac{\partial u_0^{(1)}}{\partial X} = 0, \quad (5)$$

$$\frac{\partial^2 u_0^{(1)}}{\partial X^2} = -\frac{p_0 k^2}{v_{g,L}^2 - \omega_{L,0}^2} \frac{\partial}{\partial X} |u_1^{(1)}|^2. \quad (6)$$

Here $v_{g,L} = \omega_L'(k)$, and $\{X, T\}$ are slow variables (as above).

We have defined:

$$p_0 = -r_0^3 U'''(r_0)/M \equiv 2a_{20} r_0^3, \quad q_0 = U''''(r_0) r_0^4 / (2M) \equiv 3a_{30} r_0^4$$

(both positive, and similar in magnitude for Debye interactions [4, 12]); recall that U is the interaction potential.

Eqs. (5), (6) can be combined into an NLSE in the form of Eq. (3), for $A = u_1^{(1)}$ here, with $P = P_L = \omega_L''(k)/2 < 0$. The sign of $Q > 0 (< 0)$ [11] prescribes stability (instability) at low (high) k .

Longitudinal envelope excitations are *asymmetric*: rarefactive bright or compressive dark envelope structures.




Fig. (a) Bright type; (b) dark type *asymmetric* envelope solitons.

5. Longitudinal solitons

Equation (4) is essentially the equation of atomic motion in a chain with anharmonic springs, i.e. in the celebrated FPU (*Fermi-Pasta-Ulam*) problem. At a first step, one may adopt a continuum description, viz. $\delta x_n(t) \rightarrow u(x, t)$. This leads to different nonlinear evolution equations (depending on the simplifying hypotheses adopted), some of which are critically discussed in [12]. What follows is a summary of the lengthy analysis therein.

Keeping lowest order nonlinear and dispersive terms, $u(x, t)$ obeys

$$\ddot{u} + \nu \dot{u} - c_L^2 u_{xx} - \frac{c_L^2}{12} r_0^2 u_{xxxx} = -p_0 u_x u_{xx} + q_0 (u_x)^2 u_{xx}, \quad (7)$$

where $(\cdot)_x \equiv \partial(\cdot)/\partial x$; $c_L = \omega_{L,0} r_0$; p_0 and q_0 were defined above. Assuming *near-sonic propagation* (i.e. $v \approx c_L$), and defining the relative displacement $w = u_x$, one has

$$w_\tau - a w w_\zeta + \hat{a} w^2 w_\zeta + b w_{\zeta\zeta} = 0 \quad (8)$$

(for $\nu = 0$), where

$$a = p_0/(2c_L) > 0, \quad \hat{a} = q_0/(2c_L) > 0, \quad \text{and } b = c_L r_0^2/24 > 0.$$

Following Melandso [13], various studies have relied on the *Korteweg-deVries* (KdV) equation, i.e. Eq. (8) for $\hat{a} = 0$, to gain analytical insight in the *compressive* structures observed in experiments [2]. Indeed, the KdV Eq. possesses *negative (only)*, here, since $a > 0$) supersonic pulse soliton solutions for w , implying a compressive (anti-kink) excitation for u ; the KdV soliton is thus interpreted as a density variation in the crystal, viz. $n(x, t)/n_0 \sim -\partial u/\partial x \equiv -w$. Also, the pulse width L_0 and height u_0 satisfy $u_0 L_0^2 = \text{const.}$, a feature which is confirmed by experiments [2]. However, $\hat{a} \approx 2a$ in real Debye crystals (for $\kappa \approx 1$), which invalidates the KdV approximation $\hat{a} \approx 0$ [12]). Instead, one may employ the *extended KdV* Eq. (eKdV) (8), which accounts for *both* compressive *and* rarefactive lattice excitations (exact expressions in [12]). Alternatively, Eq. (7) can be reduced to a *Generalized Boussinesq* (GBq) Equation [12]; again, for $q_0 \sim \hat{a} \approx 0$, one recovers a *Boussinesq* (Bq) equation, widely studied in solid chains. The GBq (Bq) equation yields, like its eKdV (KdV) counterpart, both compressive and rarefactive (only compressive, respectively) solutions; however, the (supersonic) propagation speed v now does *not* have to be close to c_L . The lengthy analysis (see in [12] for details) is not reproduced here.




Fig. KdV vs. Boussinesq (displacement) solitons, varying Mach no. $M = v/c_L$.

6. Longitudinal Discrete Breathers ?

Following existing studies on *Discrete Breathers* (ILMs) in *FPU chains* [cf. (4) above], it is natural to anticipate the existence of such localized excitations associated with longitudinal dust grain motion. A detailed investigation, in terms of real experimental parameters, is on the way and will be reported soon.

References

- [1] S. Takamura et al., *PoP* **8**, 1886 (2001); T. Misawa et al., *PRL* **86**, 1219 (2001); B. Liu et al., *PRL* **91**, 255003 (2003).
- [2] D. Samsonov et al., *PRL* **88**, 095004 (2002); K. Avinash et al., *PRE* **68**, 046402 (2003).
- [3] C. Zafra et al., *Phys. Rev. E* **63**, 066403 (2001); A. V. Ivlev et al., *PRL* **85**, 4060 (2000).
- [4] I. Kourakis and P. K. Shukla, *Physica Scripta* **T113**, 97 (2004).
- [5] I. Kourakis and P. K. Shukla, *Phys. Plasmas*, **11**, 2322 (2004); *idem*, **11**, 3665 (2004).
- [6] D. K. Campbell, S. Flach and Yu. S. Kivshar, *Physics Today*, **57** (1) (2004).
- [7] I. Kourakis and P. K. Shukla, *Phys. Plasmas*, **12**, 014502 (2005).
- [8] I. Kourakis, P. K. Shukla and V. Basiotis, *Proc. 2004 ICPP* (Nice, France): hal.ccsd.cnrs.fr/ccsd-00001892/en/.
- [9] V. Koukoulouyannis and I. Kourakis, *Physical Review E*, **76**, 016402/1-10 (2007).
- [10] V. Koukoulouyannis and I. Kourakis, *Multi-site breather excitations in hexagonal dusty plasma lattices*, in preparation.
- [11] I. Kourakis and P. K. Shukla, *Phys. Plasmas*, **11**, 1384 (2004).
- [12] I. Kourakis and P. K. Shukla, *Eur. Phys. J. D*, **29**, 247 (2004).
- [13] F. Melandso, *Phys. Plasmas* **3**, 3890 (1996).